

WHAT RISK/WOUND STAGE IS THE INDIVIDUAL?

HIGH RISK
AND/OR STAGE III-IV
BRADEN SCALE <11

MEDIUM RISK
AND/OR STAGE I - II
BRADEN SCALE 12-14

LOW RISK
AND/OR STAGE I
BRADEN SCALE >15

* These cushions have an available cover option utilizing GlideWear®. This patented and exclusive technology reduces shear forces measured underneath the at risk bony prominences by up to 70% thus maximizing the amount of skin protection for the prevention and treatment of pressure ulcers.

***VECTOR X**



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

***EMBRACE**



Skin Protection Positioning
Up to Wound Stage II

ASCENT



Skin Protection
Up to Wound Stage I

***ADJUSTER X**



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

***M2**



Skin Protection Positioning
Up to Wound Stage II

HYALITE



Skin Protection
Up to Wound Stage I

***VERSA X**



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

***MAXX**



Skin Protection Positioning
Up to Wound Stage II

CURVE



General Use

***LIBERTY X**



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

RIDGE



Positioning
Up to Wound Stage II

ELEMENTS W/GEL




General Use

***EMBRACE WITH ATI**




Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

***M2 WITH ATI**



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

VECTOR STANDARD & LOW PROFILE



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

ADJUSTER STANDARD & LOW PROFILE



Adjustable Medium Skin Protection Positioning
Up to Wound Stage IV

RISK / WOUND GUIDELINES

“Pressure Ulcer is a localized injury to the skin and/or underlying tissue usually over a bony prominence, as a result of pressure, or pressure in combination with shear” (NPUAP, EPUAP, PPIA, 2014).*

HIGH RISK CATEGORY/STAGE III - IV

MEDIUM RISK CATEGORY/STAGE I - II

LOW RISK CATEGORY/STAGE I

- Stage I:** Intact skin with non-blanchable redness of localized area usually over a bony prominence.
- Stage II:** Partial thickness skin loss (or blister) presenting as a shallow open ulcer with a red pink wound bed.
- Stage III:** Full thickness skin loss, subcutaneous fat may be visible but bone, tendon or muscle are not exposed.
- Stage IV:** Full thickness tissue loss with exposed bone, tendon or muscle - e.g. fascia, tendon or joint capsule. Often includes undermining and tunneling.

*National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel, Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers: Clinical Practice Guideline. Emily Haesler (Ed.). Cambridge Media: Perth, Australia; 2014.